

Influence of Nanoparticles on Seed Germination Capability, Photosynthetic Pigment Content, Activities of PSII and Superoxide Dismutase in Wheat Seedlings

Author's Details:

Gasimova F.I., Khanishova M.A., Taghiyeva K.R., Azizov I.V.✉

Institute of Molecular Biology and Biotechnologies, Azerbaijan National Academy of Sciences, Azerbaijan, AZ1073, Baku, Matbuat Avenue, 2A e-mail: fazilay@yahoo.com, Ibrahim.azizov47@gmail.com

✉ ibrahim.azizov47@gmail.com, +994 50 685 50 93

Aim

The effect of nanoparticles of copper, titanium, iron and aluminum oxides on the germination energy, germination capability, growth and development, the content of photosynthetic pigments, the activity of photosystems and superoxide dismutase of wheat seedlings was studied.

Methods. *The objects of the research were bread wheat (*Triticum aestivum* L) seedlings. The seeds of the experimental plants were treated with powders of nanoparticles of CuO, Fe₂O₃, ZnO, Al₂O₃, and TiO₂, then sown in Petri dishes and in the pots with soil.*

Results. *Nanoparticles were found to express various effects: CuO, and Al₂O₃ decreased the growth of seedlings, whereas TiO₂, ZnO and Fe₂O₃ accelerating the growth of seedlings had a positive effect on the activities of PSII and superoxide dismutase.*

Conclusion. *Nanoparticles of TiO₂, ZnO and Fe₂O₃ had a positive effect on seed germination and accelerated the growth of wheat seedlings. Nanoparticles of ZnO and Fe₂O₃ also positively influenced on the activity of PS II, and the activity of the antioxidant enzyme-superoxide dismutase also increased.*

Keywords: *wheat, nanoparticles, germination, growth of seedlings, PS II, superoxide dismutase*

Currently, nanoparticles are widely used in many branches of agriculture. It is well-known that when microelements are introduced in the form of their water-soluble salts, most of them are absorbed by the soil colloids and their absorption by plant roots is obstructed. The accumulation of these substances in the soil solution leads to environmental pollution. Nanoparticles are distinguished by unusual physicochemical properties, peculiarities of action on living organisms [1]. Thanks to the microscopic size, nanoparticles can easily pass through biological membranes, accumulate in the internal environment, and accelerate the rate of metabolic processes in cells. However, it should be noted that at high doses, nanoparticles can accumulate in plants and can later be absorbed by the human organism [2, 3, 4]. According to some authors, silver nanoparticles at low concentrations can enhance germination energy and seed germination capability, growth and development, respiration intensity and activity of enzyme systems [5-8]. It is noted that silver nanoparticles mainly accumulate in the roots of plants. TiO₂, Al₂O₃, Fe₃O₄ nanoparticles were shown to have different effects depending on their concentrations [9, 10]. In these experiments, iron nanopowders at low concentrations increased the yield and grain quality of cereals. In [11], soaking seeds with titanium, aluminum, and iron nanoparticles did not affect seed germination and plant growth. It is supposed [12] that in the process of growth and development, plants can use the surface energy of nanoparticles coming from outside, which can affect the functions of the molecular structures of the cell. Thus, the results of the researches carried out with nanoparticles are contradictory, and further researches are advisable in this direction.

The purpose of this work was to study the effect of nanoparticles of copper, titanium, iron and aluminum on germination energy, germination capability, growth and development, the content of photosynthetic pigments, the activity of photosystems and antioxidant plant enzymes.

Materials and Methods

The objects of the research were bread wheat (*Triticum aestivum* L) seedlings. The seeds of the experimental plants were treated with powders of nanoparticles of CuO, Fe₂O₃, ZnO, Al₂O₃, and TiO₂, then sown in Petri dishes and in the vegetative pots with soil. The germination energy and germination capability of seeds were determined. Then morphometric parameters, the content of chlorophyll a and b, carotenoids,

the activity of PS 2 and the enzyme superoxide dismutase were determined in the course of the plant growth. Alcohol extract of leaves was used to determine the content of pigments. The content of pigments was measured using spectrophotometer SP -2000, at 665, 649 and 440 nm. The amount of pigments was calculated using the Vernon method [13]. The activity of PS 2 was determined using the device - PAM (Photosynthesis analyzer, Germany). The activity of the superoxide dismutase enzyme was determined using the method [14]. Static data processing was performed using the program “Statistics for windows”.

Results and Discussion

Nanoparticles were found to impact differently on the germination energy and germination capability (Table 1).

Table 1. Effect of nanoparticles on germination capability and morphophysiological indices of wheat seedlings

Variants	Germination energy, %	Germination capability, %	Seedling height (cm)		F _v /F _m
			14 days	19 days	
Control	70	90	3.5 ± 0,2	11 ± 0,9	0.8
CuO	70	80	4.5 ± 0,3	12 ± 0,8	0.6
TiO ₂	85	90	6.5 ± 0,5	14 ± 0,9	0.7
ZnO	90	100	6.5 ± 0,6	14 ± 0,8	0.7
Fe ₂ O ₃	85	90	7.0 ± 0,8	15 ± 0,9	0.7
Al ₂ O ₃	60	80	4.5 ± 0,5	7,0 ± 0,7	0.6

As seen in Table 1 nanoparticles of TiO₂, ZnO and Fe₂O₃ positively affected germination energy and germination capability of seeds. These nanoparticles accelerated also growth of wheat seedlings.

Table 2. Effect of nanoparticles on photosynthetic pigment contents in wheat leaves (mg/g fresh mass)

Variants	Chlorophyll a	Chlorophyll b	a +b	a/b	Carotenoids
Control	1.8	0.55	2.35	3.27	2.2
CuO	1.4	0.46	1.86	3.04	2.9
TiO ₂	1.9	0.71	2.61	2.67	3.2
ZnO	1.9	0.69	2.59	2.75	3.5
Fe ₂ O ₃	2.1	0.72	2.82	2.92	3.7
Al ₂ O ₃	1.9	0.61	2.51	3.11	3.9

Nanoparticles of ZnO and Fe₂O₃ were also found to have a positive effect on the contents of chlorophyll a and b (Table 2).

The activity of PSII, where water photooxidation occurred with the release of hydrogen protons and molecular oxygen, increased under the influence of the nanoparticles of ZnO and Fe₂O₃. Nanoparticles of TiO₂, ZnO and Fe₂O₃ increased the activity of superoxide dismutase (Figure).




Fig. Effect of nanoparticles of biogenic metals on the activity of superoxide dismutase (in conditional units). Each bar represents mean \pm sd (standard deviation) for average of $n=3$ independent experiments; $P = 0,05$.

The results we obtained are consistent with the data obtained by Yegorov et al. [12]. They showed that pre-sowing treatment of plant seeds with low concentrations of iron nanopowder had a positive effect on the germination energy. However, an increase in the concentration by an order of magnitude led to the suppression of their growth.

Nanoparticles of TiO₂ and Al₂O₃ were found to retard the growth of bean, wheat and amaranth [10]. Studies on the effect of iron nanopowder on the growth, development, drought tolerance and productivity of maize, wheat and sunflower showed that the yield of these crops increased on average by 15-20%. The contents of gluten in the grain, oil in seeds and amino acids in the leafy stalk mass of fodder crops increased. Iron nanoparticles were shown to have different effects on the content of photosynthetic pigments in *Triticum vulgare* [15]. Nanoparticles of Fe₃O₄ had a greater effect on the total amount of photosynthetic pigments compared with those of FeO. The degree of the effect of nanoparticles on the content of photosynthetic pigments was found to depend on their concentrations.

Conclusions. Nanoparticles of TiO₂, ZnO and Fe₂O₃ had a positive effect on the germination capability and accelerated the growth of wheat seedlings. ZnO and Fe₂O₃ nanoparticles also positively influenced on the activity of PS 2, where water photooxidation occurred with the release of hydrogen protons and molecular oxygen, and increased the activity of the antioxidant enzyme superoxide dismutase.

REFERENCES

- i. Yurin V.M., Molchan O.V. *Nanomaterials and plants: a look at the problem. Works BSU. 2015. Vol.10.Part1. pp.9-21.*
- ii. R. De la Torre-Roche. *Impact of Ag nanoparticle exposure on p,p'-DDE bioaccumulation by Cucurbita pepo (zucchini) and Glycine max (soybean) / Environ. Sci. Technol. -2013. - Vol. 47. – pp. 718–725.*
- iii. 3.. Zhu H, Han J, Xiao J.Q, Jin Y. *Uptake, translocation and accumulation of manufactured iron oxide nanoparticles by pumpkin plants/ Journal Environment Monitoring. 2008. - № 10. - pp. 713–717.*
- iv. Morgalev Yu.N. *Biotesting of Nanomaterials: on the possibility of translocating nanoparticles to food networks.*
- v. Fedorenko V.F., Yerokhin M.N., Balabanov V.I., Buklagin D.S., Golubev I.G., Ishenko S.A., *Nanotechnologies and nanomaterials in the agro-industrial complex: Scientific publication. - M. : “Rosinformagrotech” FSBSI - 2011. – p.312.*
- vi. Yurkova I.N., Omelchenko A.V., Bugara I.A, *The effect of silver nanoparticles on wheat growth processes. Vestnik VSGUTU - 2014. -No1 - pp. 69–73*

- vii. Omelchenko A.V. Yurkova I.N., Jijina I.N. *The stimulating effect of silver nanoparticles on the growth and development of wheat plants: Scientific notes of Tavrichesky national university. Series of "Biology, Chemistry". - 2014 - Vol. 27 (66). - No1. - pp. 127–135.*
- viii. Kuamri, M. *Effect of silver nanoparticle (SNPs) on protein and DNA content to tomato seed (L. esculentum), cucumber (Cucumis sativus) and maize (Zea mays), / M. Kuamri, A. Mukherjee, N. Chandrasekaran : IJHGMBMS. – 2012. – V. 1, No. 1. – pp. 7–15.*
- ix. Salama, H. *Effects of silver nanoparticles in some crop plants, Common bean (Phaseolus vulgaris L.) and corn (Zea mays L.) / H. Salama // J. Biotechnology. – 2012. – V. 3, No. 10. – P. 190–197*
- x. Astafurova T.P., Morgalev Yu.N., Zotikova A.P. Verchoturova G.S., Mikhaylova S.I., Burenina A.A., Zaytseva T.A. *The effect of nanoparticles of titanium dioxide and aluminum oxide on morphophysiological parameters of plants. Bulletin of Tomsk State University. Biology-2011.-No1 (13). pp. 113-122*
- xi. Yegorov N.I., Shafronov O.D., Yegorov D.N., Suleymanov E.V. *Developing and performing an experimental evaluation of the effectiveness of application of new types of fertilizers obtained using nanotechnology in crop production. Vestnik of Nizhny Novgorod University. - 2008. - No 6. - pp. 94–99.*
- xii. Kovalenko L.V., Folmanis G.E. *Biologically active iron nanopowders / L.V. Kovalenko, - Moscow: Nauka - 2006. – p.124.*
- xiii. Shlyk A.A. *Determination of chlorophylls and carotenoids in green leaf extracts // Biochemical methods in plant physiology / M. : Nauka, 1971. pp.154-170.*
- xiv. Syrota T.V. *The method for determining the antioxidant activity of superoxide dismutase and chemical compounds. The patent of the Russian Federation No 2144674.*
- xv. Lebedev S.V., Korotkova A.M., Osipova E.A. *Evaluation of nanoparticles of Fe, Fe₃O₄ and FeSO₄ for the content of photosynthetic pigments in Triticum vulgare.// Plant Physiology. 2014, No4. pp.603-607.*